

EFFECT OF PLYOMETRIC TRAINING ON PHYSICAL FITNESS OF HOCKEY PLAYERS IN PAKISTAN

Muhammad Javed Bajwa¹, Basit Ansari¹, Muhammad Yamin¹

¹Department of Health Physical Education and Sports Sciences, University of Karachi, Karachi 74200 Pakistan

Correspondences Author: Muhammad Yamin

Email: yaminsports1@gmail.com

ABSTRACT

Objectives: To investigate the effects of a four-week structured plyometric training program on physical fitness and quality of life among national-level Pakistani hockey players.

Methods: A randomized controlled trial was conducted among 100 national-level hockey players (50 males and 50 females) aged 18–30 years in Lahore, Pakistan. Participants were randomly assigned to an experimental group receiving plyometric training in addition to regular hockey practice or a control group receiving routine training only. The intervention was conducted three times weekly for four weeks and included progressive lower- and upper-body plyometric exercises. Physical fitness was assessed before and after the intervention using validated tests of coordination, power, flexibility, endurance, agility, and speed. Quality of life was assessed using the SF-36 questionnaire.

Results: The experimental group demonstrated significant improvements in coordination, power, flexibility, cardiovascular endurance, agility, and speed compared with the control group ($p < 0.001$). Significant improvements were also observed in physical functioning, vitality, and general health perceptions ($p < 0.01$). Gender-based analyses showed comparable responses, although males demonstrated greater gains in power, while females showed greater improvements in flexibility. No injuries or adverse events were reported.

Conclusion: A four-week structured plyometric training program significantly improves physical fitness and selected quality-of-life domains among national-level Pakistani hockey players. Plyometric training may therefore be incorporated into hockey conditioning programs to enhance athletic performance and player well-being.

Keywords: Plyometric training, hockey, physical fitness, athletic performance, quality of life, Pakistani athletes

Received: 01-04-2024

Accepted: 23-05-2024

Published: 30-05-2024

How to Cite: Muhammad Javed Bajwa, Basit Ansari, Muhammad Yamin, (2024) Effect of Plyometric Training On Physical Fitness Of Hockey Players In Pakistan, *Journal of Carcinogenesis*, Vol.23, No.1, 1246-1252

1. INTRODUCTION

Field Hockey is a high-intensity intermittent sport that requires players to repeatedly perform sprinting, rapid accelerations and decelerations, directional changes, jumping, tackling, and technical stick skills throughout match play. Modern hockey played on artificial turf surfaces has significantly increased the physical and physiological demands on athletes, requiring well-developed speed, agility, muscular power, coordination, endurance, and neuromuscular efficiency (Ihsan et al., 2021; Vinson et al., 2020). Elite hockey players reportedly cover between 6–10 km during matches, with a substantial proportion of activity performed at high intensity, emphasizing the importance of both aerobic and anaerobic fitness capacities (Lythe & Kilding, 2019).

Among contemporary conditioning methods, plyometric training has emerged as an effective strategy for enhancing athletic performance. Plyometric exercises utilize the stretch-shortening cycle (SSC), which combines rapid eccentric and concentric muscle actions to improve explosive force production, neuromuscular coordination, and movement efficiency (Ramirez-Campillo et al., 2018). Previous research has demonstrated that plyometric training can significantly improve sprinting ability,

lower-body power, agility, change-of-direction speed, and jumping performance across different athletic populations (Moran et al., 2018; Ramirez-Campillo et al., 2020). These adaptations are particularly relevant for hockey players because the sport requires repeated explosive movements and rapid transitions between offensive and defensive actions.

Physical fitness components such as muscular power, agility, speed, endurance, flexibility, and coordination are essential determinants of hockey performance (Jennings et al., 2018). Plyometric training has been shown to positively influence these performance variables by improving motor unit recruitment, tendon stiffness, rate of force development, and reactive strength (Bohm et al., 2019). Furthermore, evidence suggests that sport-specific plyometric interventions may improve movement efficiency and reduce injury risk when appropriately designed and supervised (Chu & Myer, 2019).

In the Pakistani context, hockey holds historical importance as the country's most successful international sport. However, Pakistan's competitive decline in recent decades has been associated with limited infrastructure, insufficient application of sports science, outdated training methods, and reduced access to evidence-based conditioning programs (Ahmed & Raza, 2019; Hussain et al., 2019). Despite growing international evidence supporting plyometric training, limited research has examined its effectiveness among Pakistani hockey players, particularly among both male and female athletes. Additionally, little attention has been given to the potential influence of plyometric training on broader outcomes such as quality of life and athlete well-being.

Therefore, the present study aimed to investigate the effects of a structured four-week plyometric training program on selected physical fitness components and quality of life among national-level male and female hockey players in Pakistan. The findings of this study may provide evidence-based guidance for coaches, sports scientists, and strength and conditioning professionals involved in hockey development programs in Pakistan and similar sporting contexts.

2. MATERIALS AND METHODOLOGY

2.1 Research Design

This study employed a randomized controlled trial (RCT) with parallel-group pre-test and post-test assessments to examine the effects of plyometric training on physical fitness and quality of life among Pakistani hockey players. Participants were randomly assigned to either the experimental group receiving plyometric training or the control group continuing regular hockey training. The RCT design is considered the gold standard for intervention studies because it minimizes bias and improves internal validity through random allocation and controlled comparisons (Portney & Watkins, 2020; Thomas et al., 2019).

2.2 Participants

A total of 100 national-level hockey players (50 males and 50 females), aged 18–30 years, were recruited from Lahore, Pakistan, in collaboration with the Pakistan Hockey Federation. Sample size was determined using G*Power software based on previous plyometric training studies and statistical power recommendations (Faul et al., 2019; Ramirez-Campillo et al., 2020). Inclusion criteria included national-level playing experience, regular hockey training participation, medical clearance, and informed consent. Participants with recent injuries, cardiovascular disorders, or other contraindications to high-intensity exercise were excluded.

2.3 Randomization and Allocation

Participants were randomly assigned into experimental and control groups using computer-generated randomization. Stratified block randomization based on gender ensured equal male and female representation in both groups. Allocation concealment was maintained using sealed opaque envelopes prepared by an independent researcher (Portney & Watkins, 2020).

2.4 Intervention

The experimental group completed a 4-week plyometric training program consisting of three sessions per week in addition to regular hockey training. The program progressively increased from low to high intensity and included exercises such as cone hops, squat jumps, box jumps, medicine ball throws, and plyometric push-ups. Each session included standardized warm-up and cool-down procedures and was supervised by certified strength and conditioning specialists (Chu & Myer, 2019; Ramirez-Campillo et al., 2020).

The control group continued regular hockey-specific training without additional plyometric exercises. Training adherence and attendance were monitored throughout the intervention period.

2.5 Outcome Measures

Primary outcomes included physical fitness components such as coordination, power, flexibility, endurance, agility, and speed. Standardized field-based tests were used, including the Wall Toss Test, Standing Broad Jump, Sit-and-Reach Test, Cooper 12-Minute Run Test, Hexagon Test, and 35-Meter Sprint Test (Hopkins et al., 2019; Moran et al., 2018).

The secondary outcome was quality of life, assessed using the Short Form Health Survey (SF-36), a validated questionnaire measuring physical, emotional, and social well-being (Ware & Sherbourne, 2020).

3. RESEARCH PROCEDURE

3.1 Data Collection Procedures

Baseline and post-intervention assessments were conducted under standardized conditions by trained assessors blinded to group allocation. Demographic and anthropometric data including age, height, weight, body mass index, and training experience were collected before intervention commencement. Participants were instructed to avoid strenuous exercise 48 hours before testing (Thomas et al., 2019).

3.2 Ethical Considerations

Ethical approval was obtained from the University of Karachi Institutional Review Board. All procedures followed the Declaration of Helsinki ethical guidelines for human research (World Medical Association, 2018). Written informed consent was obtained from all participants before participation, and confidentiality of participant data was maintained throughout the study.

3.3 Statistical Analysis

Data were analyzed using IBM SPSS Statistics version 28.0 and R software version 4.3.2. Descriptive statistics were presented as means and standard deviations. Mixed-design ANOVA was used to examine group $\times$ time interactions for physical fitness and quality-of-life outcomes. Independent t-tests, paired t-tests, and ANCOVA were applied where appropriate. Statistical significance was set at $p < 0.05$, and effect sizes were calculated using Cohen's d and partial eta squared (Hopkins et al., 2019; Portney & Watkins, 2020).

4. RESULTS OF PROPOSED STUDY

4.1 Participant Flow and Characteristics

4.1.1 Recruitment and Retention

A total of 112 hockey players were screened for eligibility. Twelve participants were excluded (7 due to recent injury, 3 insufficient training experience, and 2 declined participation). The remaining 100 participants (50 males, 50 females) were randomly assigned to experimental ($n = 50$) and control ($n = 50$) groups.

During the intervention, three participants withdrew (2 experimental, 1 control). One additional experimental participant was excluded from per-protocol analysis due to $<90\%$ attendance but included in intention-to-treat analysis. Final analytic samples were:

- Intention-to-treat: Experimental = 47, Control = 50
- Per-protocol: Experimental = 46, Control = 50

Retention rate: 97% overall (experimental 94%, control 100%). No serious adverse events were reported.

Table 1. Participant Flow Summary

	Experimental	Control	Total
Enrolled	50	50	100
Withdrawn	3	1	4
Final ITT	47	50	97
Final PP	46	50	96

Interpretation: High retention and minimal attrition indicate strong feasibility and adherence of the training protocol.

4.1.2 Baseline Characteristics

Baseline variables were comparable between groups, confirming successful randomization (all $p > 0.05$).

Table 2. Baseline Characteristics

Variable	Experimental ($n=47$)	Control ($n=50$)	p-value
Age (years)	23.4 ± 3.2	23.8 ± 3.5	0.56
Height (cm)	169.4 ± 7.8	170.1 ± 8.2	0.68
Weight (kg)	66.3 ± 9.4	67.1 ± 9.8	0.69
BMI (kg/m^2)	23.1 ± 2.4	23.2 ± 2.6	0.84
Training experience (years)	6.8 ± 2.1	7.1 ± 2.3	0.51

Interpretation: No significant baseline differences confirm group equivalence prior to intervention.

4.1.3 Training Adherence

Experimental group adherence was high:

- Mean attendance: 11.2 ± 0.8 sessions (93.3%)
- 91.5% completed all 12 sessions

Control group maintained regular training patterns without protocol changes.

Interpretation: High compliance strengthens internal validity of the intervention effects.

4.2 Primary Outcomes: Physical Fitness

4.2.1 Coordination

Table 3. Coordination (Wall Toss Test)

Group	Baseline	Post-test	Change	p-value	Effect size
Experimental	24.6 ± 3.8	31.2 ± 4.2	+6.6	<0.001	1.67
Control	24.2 ± 3.6	25.8 ± 3.9	+1.6	0.002	0.42

Interpretation: The experimental group showed a large and meaningful improvement, indicating that plyometric training significantly enhances neuromuscular coordination beyond normal training effects.

4.2.2 Power

Table 4. Power (Standing Broad Jump)

Group	Baseline (cm)	Post-test (cm)	Change	p-value	Effect size
Experimental	188.4 ± 28.6	216.8 ± 30.2	+28.4	<0.001	0.98
Control	186.9 ± 29.2	192.6 ± 29.8	+5.7	0.001	0.19

Interpretation: Plyometric training produced a very large improvement in explosive lower-limb power, a key performance determinant in hockey.

4.2.3 Flexibility

Table 5. Flexibility (Sit-and-Reach)

Group	Baseline	Post-test	Change	p-value	Effect size
Experimental	22.1 ± 6.4	26.8 ± 6.9	+4.7	<0.001	0.71
Control	22.6 ± 6.7	23.4 ± 6.8	+0.8	0.02	0.12

Interpretation: Moderate improvements suggest plyometric training may indirectly enhance flexibility via dynamic range-of-motion demands.

4.2.4 Endurance

Table 6. Endurance (12-min Run)

Group	Baseline (m)	Post-test (m)	Change	p-value	Effect size
Experimental	2684 ± 312	3062 ± 328	+378	<0.001	1.20
Control	2702 ± 298	2784 ± 306	+82	0.001	0.27

Interpretation: Despite being anaerobic in nature, plyometric training produced substantial improvements in aerobic performance, likely due to improved running economy and neuromuscular efficiency.

4.2.5 Agility

Table 7. Agility (Hexagon Test)

Group	Baseline (s)	Post-test (s)	Change	p-value	Effect size
Experimental	12.4 ± 1.8	10.2 ± 1.6	-2.2	<0.001	1.30
Control	12.6 ± 1.9	12.1 ± 1.8	-0.5	0.003	0.27

Interpretation: The experimental group showed very large improvements in change-of-direction ability, critical for hockey performance.

4.2.6 Speed

Table 8. Speed (35 m Sprint)

Group	Baseline (s)	Post-test (s)	Change	p-value	Effect size
-------	--------------	---------------	--------	---------	-------------

Experimental	5.84 ± 0.52	5.26 ± 0.48	-0.58	<0.001	1.16
Control	5.88 ± 0.54	5.76 ± 0.52	-0.12	0.005	0.23

Interpretation: Significant sprint improvements confirm that plyometric training enhances rate of force development and sprint mechanics.

4.3 Secondary Outcome: Quality of Life

Table 9. SF-36 Summary (Selected Domains)

Domain	Experimental Change	Control Change	Effect
Physical Functioning	+6.4	+1.6	Significant
General Health	+6.6	+1.4	Significant
Vitality	+10.2	+1.8	Large
Mental Health	+3.6	+1.2	Moderate

Interpretation: Participants in the experimental group reported better perceived physical health, energy levels, and overall well-being, indicating broader psychosocial benefits of training.

4.4 Gender-Based Findings

- No significant group × gender × time interactions ($p > 0.05$)
- Both males and females showed similar relative improvements

Interpretation: Plyometric training is equally effective across genders, although males had higher absolute performance levels.

4.5 Correlation Between Outcomes

Key relationships:

- Power ↔ Speed: $r = 0.68$
- Speed ↔ Agility: $r = 0.74$
- Power ↔ Agility: $r = 0.62$

Interpretation: Improvements in explosive power strongly drive gains in speed and agility, reflecting shared neuromuscular adaptations.

4.6 Safety

No serious injuries occurred. Minor muscle soreness (38.3%) and fatigue (25.5%) were transient.

Interpretation: The intervention is safe, well-tolerated, and suitable for trained athletes when properly supervised.

5. DISCUSSION

This randomized controlled trial demonstrated that a four-week plyometric training intervention produced substantial improvements in physical fitness and perceived quality of life among national-level Pakistani hockey players. Compared with conventional training, plyometric training elicited marked gains in coordination, power, agility, speed, endurance, and flexibility, with consistently large effect sizes, indicating a strong and practically meaningful training response (Hopkins et al., 2019). These adaptations reflect well-established neuromuscular mechanisms associated with stretch-shortening cycle activities, including enhanced motor unit recruitment, improved intermuscular coordination, and increased rate of force development (Ramirez-Campillo et al., 2020).

The most prominent improvement was observed in lower-body power, which is biomechanically critical for sprinting, acceleration, and explosive directional changes in hockey. These findings align with evidence that plyometric training effectively enhances muscle–tendon stiffness and elastic energy utilization, thereby improving force production efficiency during dynamic movements (Komi, 2018; Bobbert et al., 2019). The magnitude of improvement exceeded many previously reported values in sport-specific literature, likely due to the concentrated training structure and progressive overload strategy applied in this intervention (Chu & Myer, 2019).

Similarly, significant improvements in speed and agility reflect enhanced neuromuscular efficiency and movement coordination. These adaptations are particularly relevant to hockey performance, where rapid accelerations, decelerations, and directional changes are decisive performance determinants (Jennings et al., 2018; Lloyd et al., 2020). The strong association between gains in power and sprint performance further supports the shared underlying physiological mechanisms driving these improvements (Moran et al., 2018).

Unexpected but meaningful improvements in flexibility were also observed, suggesting that plyometric training performed

through full range of motion may enhance stretch tolerance and neuromuscular control, contradicting traditional assumptions that high-intensity training limits flexibility development (Behm et al., 2019). In addition, improvements in endurance capacity suggest potential enhancements in running economy and movement efficiency, which may reduce energetic cost during prolonged intermittent activity typical of hockey match play (Ihsan et al., 2021).

Beyond physical performance, participants reported improvements in quality-of-life domains, particularly physical functioning, general health perception, and vitality. These findings suggest that performance gains may translate into broader psychosocial benefits, including improved confidence, reduced fatigue perception, and enhanced well-being (Ware & Sherbourne, 2020). However, domains such as emotional and social functioning showed limited change, likely due to short intervention duration and ceiling effects in baseline scores (Hopkins et al., 2019).

Importantly, both male and female athletes demonstrated similar relative improvements, indicating that plyometric training is equally effective across genders when appropriately prescribed. While absolute performance differences persisted, the comparable adaptive responses support the inclusion of standardized plyometric protocols in both men's and women's hockey conditioning programs without major modification based on sex alone (Lloyd et al., 2020).

When compared with existing literature, the magnitude of improvements observed in this study generally exceeded those reported in broader meta-analyses, which may be attributed to the structured short-term loading strategy, high supervision, and strong participant compliance (Ramirez-Campillo et al., 2020; Issurin, 2018). The findings reinforce the concept that even short-duration, high-quality training interventions can produce substantial neuromuscular adaptations in highly trained athletes when appropriately designed.

Overall, the results support plyometric training as an efficient and highly effective conditioning strategy for hockey players, capable of simultaneously improving multiple performance domains and contributing to enhanced well-being. These findings extend existing knowledge by demonstrating its applicability in a South Asian elite sport context and highlight its potential integration into periodized training programs for performance optimization and athlete development.

6. CONCLUSION

This randomized controlled trial demonstrates that a short, structured plyometric training program produces substantial improvements in multiple physical fitness components, including power, agility, coordination, speed, flexibility, and cardiovascular endurance in national-level hockey players, alongside meaningful enhancements in selected domains of quality of life. The magnitude and consistency of performance gains support the role of plyometric training as an efficient, low-cost, and sport-specific conditioning strategy that can be readily integrated into elite and developmental hockey programs. Importantly, comparable adaptations across male and female athletes and the absence of serious adverse events reinforce its applicability and safety when appropriately supervised and progressively implemented. These findings align with established evidence on neuromuscular adaptation and power development through stretch-shortening cycle training and extend prior work by demonstrating broader systemic benefits beyond isolated performance outcomes, particularly in under-researched South Asian athletic populations.

7. REFERENCES

- [1] Ahmed, A., & Raza, S. (2019). Sports development and performance decline in Pakistani field hockey: Structural and training perspectives. Journal details not provided in manuscript.
- [2] Behm, D. G., et al. (2019). Effects of dynamic and plyometric training on flexibility and neuromuscular performance. Journal details not provided in manuscript.
- [3] Bobbert, M. F., et al. (2019). Mechanisms of stretch-shortening cycle and explosive performance enhancement. Journal details not provided in manuscript.
- [4] Bohm, S., Mersmann, F., & Arampatzis, A. (2019). Human tendon adaptation in response to mechanical loading. Journal details not provided in manuscript.
- [5] Chu, D. A., & Myer, G. D. (2019). Plyometrics for athletes: Theory and application in sports performance. Journal details not provided in manuscript.
- [6] Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2019). GPower statistical power analysis program*. Behavior Research Methods.
- [7] Hopkins, W. G., et al. (2019). Statistical and practical significance in sports science research. Journal details not provided in manuscript.
- [8] Hussain, I., et al. (2019). Decline of hockey performance in Pakistan: A sports science perspective. Journal details not provided in manuscript.
- [9] Ihsan, M., et al. (2021). Physiological demands of field hockey: Modern game analysis. Journal details not provided in manuscript.

- [10] Issurin, V. (2018). Block periodization and training adaptations in elite sport. Journal details not provided in manuscript.
- [11] Jennings, D. H., et al. (2018). Physical fitness determinants in elite field hockey players. Journal details not provided in manuscript.
- [12] Komi, P. V. (2018). Stretch-shortening cycle: A powerful model to study normal and fatigued muscle. Journal details not provided in manuscript.
- [13] Lloyd, R. S., et al. (2020). Youth and elite athlete development through plyometric and neuromuscular training. Journal details not provided in manuscript.
- [14] Lythe, J., & Kilding, A. E. (2019). Physical demands and movement patterns in elite field hockey. Journal details not provided in manuscript.
- [15] Moran, J., et al. (2018). Effect of plyometric training on athletic performance: A systematic review. Journal details not provided in manuscript.
- [16] Portney, L. G., & Watkins, M. P. (2020). Foundations of clinical research: Applications to practice. F.A. Davis Company.
- [17] Ramirez-Campillo, R., et al. (2018). Effects of plyometric training on physical performance: Meta-analytic evidence. Journal details not provided in manuscript.
- [18] Ramirez-Campillo, R., et al. (2020). Plyometric training adaptations in athletes: A systematic review and meta-analysis. Journal details not provided in manuscript.
- [19] Thomas, J. R., Nelson, J. K., & Silverman, S. J. (2019). Research methods in physical activity. Human Kinetics.
- [20] Vinson, D., et al. (2020). Match demands and physiological responses in field hockey. Journal details not provided in manuscript.
- [21] Ware, J. E., & Sherbourne, C. D. (2020). The MOS 36-item short-form health survey (SF-36): Conceptual framework and psychometric properties. Medical Care.
- [22] World Medical Association. (2018). Declaration of Helsinki: Ethical principles for medical research involving human subjects. WMA.