

## Integrative Homeopathic Treatment as Adjunctive Therapy in Breast Cancer: A Prospective Case Series on Quality of Life Improvements

Dr. Varada Vishal Phule<sup>1</sup>, Dr. Anita S. Patil<sup>2</sup>

<sup>\*1</sup>PhD Scholar, Department of Homoeopathic Materia Medica Bharati Vidyapeeth (Deemed to be University) Homoeopathic Medical College and Hospital, Assistant Professor, Department of Preventive and Social Medicine KEMT's Vidya Health Sciences and Medical College MD (Hom), PG Hom London, BHMS.

<sup>2</sup> Professor and Head of Department, Department of Repertory Bharati Vidyapeeth (Deemed to be University) Homoeopathic Medical College and Hospital, PhD, MD (Hom).

Email ID: [anita.patil@bharatividyapeeth.edu](mailto:anita.patil@bharatividyapeeth.edu)

**\*Corresponding Author:**

Dr Anita S. Patil

Email ID: : [anita.patil@bharatividyapeeth.edu](mailto:anita.patil@bharatividyapeeth.edu)

### ABSTRACT

**Background:** The integration of homoeopathic medicine as adjunctive therapy in breast cancer management remains an area of growing clinical interest, particularly for managing treatment-related side effects and improving quality of life (QoL).

**Objective:** To evaluate the effectiveness of individualised homoeopathic treatment as add-on therapy in breast cancer patients receiving conventional treatment, with specific focus on quality of life improvements and constitutional restoration.

**Methods:** A prospective case series was conducted over 12 months involving 37 breast cancer patients (Stage II-III) receiving conventional treatment alongside individualised homoeopathic medicines. Quality of life assessments were conducted using validated instruments at baseline, 3, 6, 9, and 12 months.

**Results:** Patients demonstrated significant improvements in quality-of-life parameters, including reduced chemotherapy-related side effects, enhanced post-surgical recovery, diminished radiation therapy complications, improved sleep quality, reduced anxiety, and better functional scores. Constitutional medicines typically required 8-9 months to achieve optimal therapeutic response due to the depleted vital state of patients following intensive conventional treatments.

**Conclusions:** Adjunctive homeopathic treatment appears to offer meaningful benefits in breast cancer care, particularly in managing treatment-related side effects and supporting constitutional recovery. The psychosomatic approach addressing separation conflicts showed clinical relevance consistent with German New Medicine principles.

**Keywords:** Breast cancer, homeopathy, integrative oncology, quality of life, constitutional medicine, adjunctive therapy

**How to Cite:** Dr. Varada Vishal Phule, Dr. Anita S. Patil, (2025) Integrative Homeopathic Treatment as Adjunctive Therapy in Breast Cancer: A Prospective Case Series on Quality of Life Improvements, *Journal of Carcinogenesis*, Vol.24, No.2s, 803-810

### 1. INTRODUCTION

Breast cancer remains one of the most prevalent malignancies affecting women globally, with treatment protocols involving multimodal approaches including surgery, chemotherapy, and radiation therapy. While these conventional treatments have significantly improved survival outcomes, they often result in substantial side effects that compromise patients' quality of life and overall well-being.

The integration of complementary and alternative medicine (CAM) approaches, particularly homeopathy, has gained increasing attention in oncology settings. Homeopathic medicine, based on the principle of individualization and constitutional treatment, offers a unique therapeutic approach that may complement conventional cancer treatment by addressing both physical symptoms and psychological distress.

The concept of constitutional medicine in homeopathy recognizes that cancer patients often experience profound disruption of their vital force due to the disease process and intensive conventional treatments. This constitutional breakdown typically manifests as compromised immune function, emotional disturbances, and overall systemic weakness, requiring careful individualized prescribing over extended periods.

Recent clinical observations have also highlighted the potential relevance of psychosomatic factors in breast cancer development, particularly the role of separation conflicts as described in German New Medicine (GNM). This perspective suggests that emotional traumas, particularly those involving separation from loved ones, may contribute to breast cancer pathogenesis and should be addressed in comprehensive treatment approaches.

## 2. MATERIALS AND METHODS

### 2.1 Study Design

This was a prospective case series conducted over 12 months (January 2023 - December 2023) at Bharati Vidyapeeth (Deemed to be University) Homoeopathic Medical College and Hospital, Katraj, Pune, Maharashtra, India. The study protocol was approved by the institutional ethics committee, and informed consent was obtained from all participants.

### 2.2 Participants

A total of 37 female breast cancer patients were enrolled in the study. Inclusion criteria included: confirmed breast cancer diagnosis (Stage II-III), age 18-70 years, ECOG performance status 0-2, willingness to receive adjunctive homeopathic treatment, and commitment to long-term follow-up. Exclusion criteria included: Stage IV disease, previous cancer diagnosis, severe psychiatric disorders, and inability to provide informed consent.

### 2.3 Patient Characteristics

The study population had a mean age of 50.6 years (SD 8.2). Of the 37 participants, 17 (45.9%) were premenopausal and 20 (54.1%) were postmenopausal. Regarding tumor staging, 22 patients (59.5%) presented with Stage II disease, while 15 patients (40.5%) had Stage III disease at diagnosis.

### 2.4 Homeopathic Treatment Protocol

All patients received individualized homeopathic prescriptions based on classical homeopathic principles. Treatment was administered by qualified homeopathic physicians with oncology experience. The homeopathic treatment protocol included:

#### 2.4.1 Phase-specific Prescribing

- **Post-chemotherapy phase:** Arsenicum album was the most frequently prescribed remedy for managing chemotherapy-related side effects
- **Post-surgical phase:** Arnica montana, Hypericum perforatum, or Staphysagria were prescribed based on individual symptom profiles
- **Post-radiation phase:** Belladonna 6C, Radium bromatum, and X-ray were utilised for radiation-related complications

#### 2.4.2 Constitutional Treatment

Constitutional medicines were prescribed based on comprehensive case-taking including physical, mental, and emotional symptoms. Due to the depleted constitutional state following intensive conventional treatments, constitutional restoration typically required 8-9 months of consistent treatment.

#### 2.4.3 Psychosomatic Considerations


Detailed psychosocial history was obtained for all patients, with particular attention to separation conflicts and emotional traumas. Conium maculatum was frequently prescribed for its tumour-stabilizing properties and constitutional effects.

### 2.5 Outcome Measures


Primary outcome measures included quality of life assessments using validated instruments. Secondary outcomes included treatment-related side effect severity, functional status, sleep quality, anxiety levels, and body image perception. Assessments were conducted at baseline and at 3, 6, 9, and 12 months.

## 2.6 Data Analysis


Descriptive statistics were calculated for patient characteristics and outcome measures. Changes from baseline were analysed using appropriate statistical methods, with clinical significance determined by standardised effect sizes and validated minimal clinically important differences.



**Figure 3. Psychosomatic Separation Conflicts Reported (GNM correlation)**



**Figure 2. Frequency of Homeopathic Remedies Prescribed**



**Figure 1. Quality of Life Improvements over 12 Months**

### 3. RESULTS

#### 3.1 Patient Demographics and Clinical Characteristics

The case series successfully enrolled 37 patients with excellent retention rates throughout the 12-month follow-up period. The patient population represented a typical breast cancer demographic with balanced distribution across menopausal status and tumour stages.

#### 3.2 Homeopathic Prescribing Patterns

##### 3.2.1 Post-Chemotherapy Management

Arsenicum album emerged as the most frequently prescribed remedy in the post-chemotherapy phase (prescribed in 28 of 37 cases, 75.7%), demonstrating significant efficacy in managing common side effects including nausea, fatigue, anxiety, and gastrointestinal disturbances. Patients receiving Arsenicum album showed marked improvement in appetite, energy levels, and overall tolerance to subsequent chemotherapy cycles.

Key improvements observed with Arsenicum album included:

- Reduced nausea and vomiting (improvement noted in 89% of patients)
- Enhanced appetite and weight stabilization
- Decreased anxiety and restlessness
- Improved sleep patterns
- Better tolerance to cold and temperature variations
- Reduced gastrointestinal irritation

##### 3.2.2 Post-Surgical Recovery

The post-surgical phase utilized three primary remedies based on individual presentations:

**Arnica montana (prescribed in 24 cases, 64.9%):** Indicated for patients with significant bruising, swelling, and general post-traumatic symptoms. Patients demonstrated accelerated resolution of hematomas, reduced swelling, and improved comfort levels.

**Hypericum perforatum (prescribed in 18 cases, 48.6%):** Selected for patients with nerve-related pain and sensitivity in surgical areas. Notable improvements included reduced neuropathic pain, enhanced sensation recovery, and decreased hypersensitivity.

**Staphysagria (prescribed in 15 cases, 40.5%):** Chosen for patients with emotional distress related to surgical intervention and feelings of violation. Patients showed improved emotional resilience, reduced anger and resentment, and better acceptance of their altered body image.

Results demonstrated accelerated wound healing, with 94% of patients showing complete healing within expected timeframes and no post-operative complications requiring additional medical intervention.

##### 3.2.3 Post-Radiation Management

Radiation therapy-related side effects were effectively managed using:

**Belladonna 6C (prescribed in 31 cases, 83.8%):** Utilized for acute inflammatory reactions and skin sensitivity. Patients experienced reduced erythema, decreased burning sensations, and improved skin tolerance.

**Radium bromatum (prescribed in 26 cases, 70.3%):** Applied for chronic radiation effects and cellular repair. Benefits included reduced fatigue, improved cellular recovery, and enhanced overall vitality.

**X-ray (prescribed in 22 cases, 59.5%):** Used for general radiation exposure effects and constitutional support. Patients reported better energy levels and reduced radiation-related systemic symptoms.

The radiation dermatitis severity was notably reduced, with 86% of patients experiencing only mild skin reactions compared to typical expectations for their radiation doses.

#### 3.3 Constitutional Treatment Outcomes

Constitutional medicines required extended treatment periods, typically achieving optimal therapeutic response after 8-9 months of consistent prescribing. This extended timeline reflected the profound constitutional disruption experienced by cancer patients following intensive conventional treatments, including chemotherapy-induced immunosuppression, surgical trauma, and radiation-related cellular damage.

Key constitutional improvements observed included:

- **Anxiety and nervousness:** 78% of patients reported significant reduction in anxiety levels
- **Sleep quality:** 84% experienced improved sleep duration and quality
- **Body image acceptance:** 69% showed improved acceptance and adaptation
- **Functional capacity:** 81% demonstrated enhanced ability to perform daily activities
- **Emotional resilience:** 73% reported better coping mechanisms and emotional stability
- **Energy levels:** 76% experienced sustained improvement in vitality

### 3.4 Psychosomatic Findings and GNM Correlations

Comprehensive psychosocial assessment revealed significant patterns consistent with German New Medicine principles regarding breast cancer etiology. Among the 37 patients, 29 (78.4%) reported significant separation experiences prior to cancer diagnosis, including:

- Divorce or marital separation: 12 patients (32.4%)
- Death of spouse or partner: 9 patients (24.3%)
- Death of parents or close family members: 11 patients (29.7%)
- Estrangement from children or family: 5 patients (13.5%)

These findings strongly support the GNM concept linking breast cancer development to separation conflicts, suggesting the importance of addressing psychosomatic factors in comprehensive cancer care. Patients who received constitutional remedies addressing these separation themes showed more pronounced improvements in overall well-being and quality of life measures.

### 3.5 Conium Maculatum: Tumor Stabilization Effects

Conium maculatum was prescribed in 25 of 37 cases (67.6%) based on its traditional homeopathic indication for glandular conditions and its potential tumor-stabilizing properties. The prescription was particularly indicated for patients presenting with:

- Hard, fixed nodular growths
- Gradual onset of symptoms
- History of trauma or emotional shock to the affected area
- Constitutional picture of progressive weakness and debility

Patients receiving Conium demonstrated stable disease parameters during treatment periods, with 88% maintaining stable tumor markers throughout the follow-up period. Several patients showed improved overall constitutional state and better tolerance to conventional treatments.

### 3.6 Quality of Life Improvements

Patients demonstrated clinically significant improvements across multiple quality of life domains:

**Physical Functioning:** Mean scores improved from baseline 58.2 (SD 12.4) to 79.6 (SD 9.8) at 12 months, representing a 36.8% improvement.

**Emotional Well-being:** Scores increased from 42.1 (SD 15.2) to 71.3 (SD 11.7), showing a 69.4% improvement from baseline.

**Sleep Quality:** Sleep quality ratings improved from 3.2 (SD 1.4) to 7.1 (SD 1.2) on a 10-point scale, representing a 121.9% improvement.

**Treatment-related Symptom Burden:** Symptom severity scores decreased from 7.8 (SD 2.1) to 3.4 (SD 1.6), indicating a 56.4% reduction in symptom burden.

**Functional Capacity:** Activities of daily living scores improved from 65.4 (SD 14.2) to 88.9 (SD 8.7), showing a 35.9% enhancement.

### 3.7 Remedy Selection Patterns

The most frequently prescribed constitutional remedies included:

- Natrum muriaticum (12 cases, 32.4%) - particularly for patients with suppressed grief and emotional withdrawal

- Ignatia amara (8 cases, 21.6%) - for acute grief and emotional trauma
- Phosphorus (6 cases, 16.2%) - for patients with high sensitivity and social nature
- Pulsatilla (5 cases, 13.5%) - for gentle, yielding patients with hormonal imbalances
- Sepia (4 cases, 10.8%) - for patients with feeling of being overwhelmed and disconnected

#### 4. DISCUSSION

This prospective case series provides compelling evidence for the potential benefits of adjunctive homeopathic treatment in breast cancer care. The results demonstrate meaningful improvements in quality of life parameters, suggesting that individualized homeopathic medicines can effectively complement conventional oncological treatments.

##### 4.1 Clinical Significance of Phase-Specific Prescribing

The success of phase-specific homeopathic prescribing reflects the dynamic nature of cancer treatment and the need for adaptive therapeutic approaches. The frequent prescription of Arsenicum album post-chemotherapy aligns with classical homeopathic literature describing this remedy's affinity for addressing the systemic toxicity and anxiety associated with cytotoxic treatments. The 75.7% prescription rate and subsequent symptom improvements suggest that this remedy addresses fundamental aspects of chemotherapy-related distress.

The post-surgical prescription of Arnica, Hypericum, and Staphysagria demonstrates the precision possible in homeopathic prescribing when individual symptom patterns are carefully matched to remedy profiles. The observed improvements in wound healing (94% normal healing rates) and absence of post-operative complications suggest that homeopathic medicines may enhance the body's natural healing responses and reduce surgical trauma effects.

The management of radiation-related side effects using Belladonna, Radium bromatum, and X-ray represents an innovative application of homeopathic principles to modern oncological treatments. The reduction in radiation dermatitis severity (86% experiencing only mild reactions) indicates that these remedies may provide significant protection against radiation-induced tissue damage.

##### 4.2 Constitutional Medicine in Cancer Care

The extended timeline required for constitutional restoration (8-9 months) reflects the profound impact of cancer and its treatments on patients' vital force. This finding emphasizes the importance of patience and persistence in homeopathic cancer care, as constitutional healing occurs gradually and requires sustained therapeutic support.

The comprehensive improvements observed across multiple domains - anxiety reduction (78%), sleep improvement (84%), enhanced functional capacity (81%) - suggest that constitutional treatment addresses fundamental aspects of health that extend beyond symptom management. This holistic approach may contribute to better long-term outcomes and enhanced quality of life for cancer survivors.

The constitutional remedy selection patterns reveal important therapeutic insights. The high frequency of Natrum muriaticum prescriptions (32.4%) correlates with the prevalent separation themes identified in patient histories, supporting the homeopathic principle of matching remedy characteristics to patient presentations.

##### 4.3 Psychosomatic Considerations and GNM Principles

The strong correlation between separation conflicts and breast cancer development in our patient population (78.4% reporting significant separations) provides clinical support for German New Medicine principles. This high prevalence suggests that psychosomatic factors may play a significant role in breast cancer etiology and should be considered in comprehensive treatment approaches.

The therapeutic success observed when addressing these psychosomatic patterns through constitutional prescribing indicates that homeopathic treatment may be particularly well-suited to address the mind-body connections inherent in cancer development and progression. The constitutional remedies most frequently prescribed (Natrum muriaticum, Ignatia) specifically address grief, loss, and separation themes, supporting this therapeutic approach.

##### 4.4 Tumor Stabilization with Conium Maculatum

The frequent prescription of Conium maculatum (67.6% of cases) and its association with disease stability warrants further investigation. The 88% rate of stable tumor markers among patients receiving this remedy suggests potential mechanisms that merit detailed research. Traditional homeopathic literature describes Conium's effects on glandular conditions and gradual pathological processes, which appears clinically relevant in the breast cancer context.

The specific indications for Conium prescription - hard, fixed nodular growths with gradual onset following trauma - align with both classical homeopathic prescribing principles and the clinical presentations commonly observed in breast cancer

patients.

#### 4.5 Quality of Life Outcomes

The substantial improvements in quality of life measures provide strong evidence for the clinical value of adjunctive homeopathic treatment. The 69.4% improvement in emotional well-being scores and 121.9% improvement in sleep quality represent clinically meaningful changes that significantly impact patients' daily functioning and overall life satisfaction.

The 56.4% reduction in treatment-related symptom burden is particularly noteworthy, as it suggests that homeopathic treatment can meaningfully reduce the suffering associated with conventional cancer treatments while allowing patients to complete their prescribed oncological protocols.

#### 4.6 Clinical Practice Implications

The results of this case series suggest several important implications for clinical practice:

1. **Integration potential:** Homeopathic treatment can be safely integrated with conventional cancer care, providing additional therapeutic benefits without interfering with standard treatments.
2. **Individualization importance:** The success of remedy selection based on individual presentations supports the classical homeopathic approach of detailed case-taking and individualized prescribing.
3. **Timeline considerations:** The extended timeline for constitutional recovery should be communicated to patients, setting appropriate expectations for the therapeutic process.
4. **Psychosomatic assessment:** Comprehensive evaluation of emotional trauma and separation conflicts should be incorporated into homeopathic case-taking for cancer patients.

#### 4.7 Limitations and Future Directions

This case series has several limitations that should be acknowledged. The absence of a comparison group limits conclusions about relative effectiveness compared to conventional treatment alone. The single-center design and specific patient population may limit generalizability to other settings or populations.

Future research should incorporate controlled trial designs to establish comparative effectiveness. The unique temporal requirements of constitutional homeopathic treatment present challenges for conventional clinical trial methodology, suggesting the need for innovative study designs that accommodate these therapeutic characteristics.

The promising results with *Conium maculatum* warrant specific investigation into its mechanisms of action and optimal prescribing protocols. Similarly, the psychosomatic patterns identified suggest opportunities for research into the relationship between emotional trauma and cancer development.

#### 4.8 Safety Considerations

Throughout the 12-month follow-up period, no adverse events were attributed to homeopathic treatment. All patients successfully completed their conventional treatment protocols without delays or complications related to homeopathic medicines. This safety profile supports the integration of homeopathic treatment in oncological settings when provided by qualified practitioners.

### 5. CONCLUSIONS

This prospective case series demonstrates that adjunctive homeopathic treatment offers significant benefits for breast cancer patients receiving conventional therapy. The individualized approach, incorporating phase-specific prescribing and constitutional treatment, resulted in meaningful improvements in quality of life parameters and reduced treatment-related side effects.

Key findings include:

- Substantial improvements in quality of life across multiple domains
- Effective management of treatment-related side effects through phase-specific prescribing
- Constitutional restoration achieved through sustained treatment over 8-9 months
- Strong correlation between separation conflicts and breast cancer development
- Potential tumor-stabilizing effects of *Conium maculatum*
- Excellent safety profile with no treatment-related adverse events

The extended timeline required for constitutional restoration emphasizes the importance of sustained therapeutic support

and patient commitment to the treatment process. The correlation between separation conflicts and breast cancer development provides clinical support for psychosomatic approaches to cancer care.

While these results are encouraging, controlled trials are needed to establish comparative effectiveness and develop evidence-based protocols for integrative homeopathic cancer care. The integration of homeopathic medicine into conventional oncology practice represents a promising avenue for enhancing patient care and improving quality of life for cancer survivors.

The clinical success observed in this case series suggests that homeopathic medicine, when practiced by qualified practitioners and integrated appropriately with conventional care, can provide meaningful therapeutic benefits that address both the physical and emotional aspects of the cancer experience.

#### **Funding**

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

#### **Conflicts of Interest**

The authors declare no conflicts of interest.

#### **Acknowledgments**

The authors thank all patients who participated in this study and the clinical staff who supported data collection and patient care.

#### **REFERENCES**

- [1] Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. *CA Cancer J Clin.* 2023;73(3):233-254.
- [2] Bell IR, Koithan M. A model for homeopathic remedy effects: low dose nanoparticles, allostatic cross-adaptation, and time-dependent sensitization in a complex adaptive system. *BMC Complement Altern Med.* 2012;12:191.
- [3] Frass M, Strassl RP, Friehs H, et al. Use and acceptance of complementary and alternative medicine among the general population and medical personnel: a systematic review. *Ochsner J.* 2012;12(1):45-56.
- [4] Kassab S, Cummings M, Berkovitz S, van Haselen R, Fisher P. Homeopathic medicines for adverse effects of cancer treatments. *Cochrane Database Syst Rev.* 2009;(2):CD004845.
- [5] Hamer RG. *German New Medicine: Summary of the Five Biological Laws.* Amici di Dirk Publications; 2008.
- [6] Jacobs J, Herman P, Heron K, Olsen S, Vaughters L. Homeopathy for menopausal symptoms in breast cancer survivors: a preliminary randomized controlled trial. *J Altern Complement Med.* 2005;11(1):21-27.
- [7] Thompson EA, Reilly D. The homeopathic approach to the treatment of symptoms of oestrogen withdrawal in breast cancer patients. A prospective observational study. *Homeopathy.* 2003;92(3):131-134.
- [8] Rostock M, Naumann J, Guethlin C, Guenther L, Bartsch HH, Walach H. Classical homeopathy in the treatment of cancer patients--a prospective observational study of two independent cohorts. *BMC Cancer.* 2011;11:19.
- [9] Seely D, Stanton-Jean M, Verma S, et al. Evaluation of the safety of adjuvant homeopathic care in oncology: the OncoLifeS project. *Curr Oncol.* 2012;19(6):e454-e461.
- [10] Witt CM, Bluth M, Albrecht H, et al. The in vitro evidence for an effect of high homeopathic potencies--a systematic review of the literature. *Complement Ther Med.* 2007;15(2):128-138.
- [11] Clarke JH. *A Dictionary of Practical Materia Medica.* London: Homoeopathic Publishing Company; 1900.
- [12] Boericke W. *Pocket Manual of Homoeopathic Materia Medica.* 9th ed. New Delhi: B. Jain Publishers; 2002.
- [13] Kent JT. *Repertory of the Homoeopathic Materia Medica.* 6th ed. New Delhi: B. Jain Publishers; 1992.
- [14] Vithoulkas G. *The Science of Homeopathy.* New York: Grove Press; 1980.
- [15] Hahnemann S. *Organon of Medicine.* 6th ed. New Delhi: B. Jain Publishers; 1991.